

There are some known errors in this print edition of *BNF for Children 2015–2016*.
Electronic versions have been updated

Aciclovir (pg 367)	Herpes simplex encephalitis: treatment should be high dose IV
Co-amoxiclav 400/57 suspension (pg 313)	Dose for 6-year olds omitted
Erythromycin (pg 300)	Dose for prophylaxis in asplenia or sickle-cell disease for 2–7 year olds incorrect Dose for acne for 1–2 year olds incorrect
Esomeprazole (pg 50)	Age range for IV treatment of GORD incorrect
Magnesium glycerophosphate (pg 534)	Dose for a child 1 month to 11 years should be initially 50 mg/kg 3 times a day
Paraldehyde (pg 261)	Dose based on use of a premixed solution of paraldehyde in olive oil in equal volumes
Pyrimethamine (pg 361)	Age range for congenital toxoplasmosis incorrect
Tranexamic acid (pg 72)	Dose frequency for IV treatment of fibrinolysis and max. dose for prevention of excessive bleeding incorrect
Vecuronium (pg 736)	Administration section: IV concentration incorrect

Users are reminded:

- to use digital versions at www.medicinescomplete.com or the BNF App for the most up-to-date information, including corrections to the affected monographs;
- see www.bnf.org to remain aware of clarifications, corrections, and updates.